

Datasheet for ABIN894626

anti-DVL1 antibody (AA 21-100) (AbBy Fluor® 488)



Go to Product page

Background:

Quantity:	100 μL
Target:	DVL1
Binding Specificity:	AA 21-100
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DVL1 antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human DVL1
Isotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.
Target Details	
Target:	DVL1
Alternative Name:	DVL1 (DVL1 Products)

Synonyms: DVL, DVL1L1, DVL1P1, Segment polarity protein dishevelled homolog DVL-1,

Expiry Date:

l arget Details	
	Dishevelled-1, DSH homolog 1, DVL1
	Background: Participates in Wnt signaling by binding to the cytoplasmic C-terminus of frizzled
	family members and transducing the Wnt signal to down-stream effectors. Plays a role both in
	canonical and non-canonical Wnt signaling. Plays a role in the signal transduction pathways
	mediated by multiple Wnt genes. Required for LEF1 activation upon WNT1 and WNT3A
	signaling. DVL1 and PAK1 form a ternary complex with MUSK which is important for MUSK-
	dependent regulation of AChR clustering during the formation of the neuromuscular junction
	(NMJ).
Gene ID:	1855
UniProt:	014640
Pathways:	WNT Signaling, Synaptic Membrane, Skeletal Muscle Fiber Development
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

12 months