# antibodies -online.com





## anti-EID3 antibody (AA 65-160) (HRP)



#### Overview

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | EID3  |
| Binding Specificity: | AA 65-160   |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This EID3 antibody is conjugated to HRP   |
| Application:         | Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human EID3 |
|-----------------------|--|
| Isotype:              | IgG  |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Horse                      |
| Purification:         | Purified by Protein A.                                   |

### **Target Details**

| Target:           | EID3  |
|-------------------|---|
| Alternative Name: | EID3 (EID3 Products)  |
| Background:       | Synonyms: E1A-like inhibitor of dferentiation 3, EID 3, EID-1-like inhibitor of dferentiation 3, EID- |

| Target Details      |  |
|---------------------|--|
|                     | 3, EID3_HUMAN, EP300 interacting inhibitor of dferentiation 3, EP300-interacting inhibitor of dferentiation 3, Non-SMC element 4, S. cerevisiae, homolog of, B, NSE4B, NSMCE4B.  Background: Act as a repressor of nuclear receptor-dependent transcription possibly by interfering with CREBBP-dependent coactivation. May function as a coinhibitor of other CREBBP/EP300-dependent transcription factors. Tissue specificity: Highly expressed in testis. |
| Gene ID:            | 493861   |
| Application Details |  |
| Application Notes:  | WB 1:300-5000<br>IHC-P 1:200-400<br>IHC-F 1:100-500  |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Concentration:      | 1 μg/μL  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.   |

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Handling Advice:   | Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.               |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |