# antibodies -online.com





# anti-UBR1 antibody (AA 801-900) (Alexa Fluor 647)



Go to Product page

$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	UBR1	
Binding Specificity:	AA 801-900	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This UBR1 antibody is conjugated to Alexa Fluor 647	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human eIF3A
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	UBR1
Alternative Name:	e3A (UBR1 Products)

## **Target Details**

Background:	Synonyms: EIF3, P167, p180, p185, TIF32, EIF3S10, eIF3-p170, eIF3-theta, Eukaryotic	
	translation initiation factor 3 subunit A, eIF3a, Eukaryotic translation initiation factor 3 subunit	
	10, eIF-3-theta, eIF3 p167, eIF3 p180, eIF3 p185, KIAA0139	
	Background: Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which	
	is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates	
	with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-	
	tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates	
	mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3	
	complex is also required for disassembly and recycling of post-termination ribosomal	
	complexes and subsequently prevents premature joining of the 40S and 60S ribosomal	
	subunits prior to initiation.	
Gene ID:	8661	
UniProt:	Q14152	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	