# antibodies .- online.com





# anti-ELAC2 antibody (AA 88-190) (AbBy Fluor® 555)



Go to Product page

( )	11/0	K\ /	iew
	$\cup$	'I V/I	$I \cap VV$

Quantity:	100 μL	
Target:	ELAC2	
Binding Specificity:	AA 88-190	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ELAC2 antibody is conjugated to AbBy Fluor® 555	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ELAC2
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	ELAC2
Alternative Name:	ELAC2 (ELAC2 Products)

## **Target Details**

Background:	Synonyms: elaC homolog 2, ElaC homolog protein 2, ELC2, Heredity prostate cancer protein 2,	
	HPC2, putative prostate cancer susceptibility protein HPC2/ELAC2, Ribonuclease Z2, RNase Z2,	
	tRNA 3 endonuclease 2, tRNase Z long form, tRNase Z2, Zinc phosphodiesterase ELAC protein	
	2.	
	Background: ELAC2 is a zinc phosphodiesterase which displays some tRNA 3' processing	
	endonuclease activity. It is thought to be involved in tRNA maturation, acting by removing a 3'	
	trailer from precursor tRNA. ELAC2 also interacts with activated Smad family member 2	
	(Smad2) and its nuclear partner forkhead box H1 (also known as FAST-1), and reduced	
	expression can suppress transforming growth factor-beta induced growth arrest. Mutations in	
	this gene result in an increased risk of prostate cancer. A number of allelic variants have been	
	investigated in hereditary prostate cancer.	
Gene ID:	60528	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	