

Datasheet for ABIN895753

anti-Endothelin 1 antibody (AA 53-73) (AbBy Fluor® 488)



Overview

Quantity:	100 μL
Target:	Endothelin 1 (EDN1)
Binding Specificity:	AA 53-73
Reactivity:	Human, Rat, Pig, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Endothelin 1 antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human human Endothelin-1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Pig, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Horse,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	Endothelin 1 (EDN1)
Alternative Name:	Endothelin-1 (EDN1 Products)

Target Details

_	
Background:	Synonyms: Endothelin-1, EDN 1, EDN1, EDN1 protein, Endothelin1, ET 1,ET1,
	Background: Endothelins (ET) show potent constrictor activity in vascular and non-vascular
	smooth muscle. This family of 21-amino acid peptides exists in at least three isoforms - ET-1,
	ET-2, and ET-3, and is produced in endothelial and epithelial cells. ET's can mediate biological
	effects in cells and tissues, and have been shown to bind to an ET receptor in the lung, kidney,
	heart, and liver. Endothelin 1 is expressed in lung, placental stem villi vessels and in cultured
	placental vascular smooth muscle cells.
Gene ID:	1906
Pathways:	Hormone Transport, Negative Regulation of Hormone Secretion, Regulation of Systemic Arterial
	Blood Pressure by Hormones, cAMP Metabolic Process, Regulation of Muscle Cell
	Differentiation, Regulation of G-Protein Coupled Receptor Protein Signaling, Regulation of Cell
	Size
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months