antibodies -online.com





anti-FAM134C antibody (AA 76-180) (Cy5)



| () | 1/0 | r\ / I | 014 | |
|-----|-----|--------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | FAM134C | |
| Binding Specificity: | AA 76-180 | |
| Reactivity: | Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This FAM134C antibody is conjugated to Cy5 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human FAM134C |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit,Guinea Pig |
| Purification: | Purified by Protein A. |

Target Details

| Target: | FAM134C |
|-------------------|----------------------------|
| Alternative Name: | FAM134C (FAM134C Products) |

Target Details

| Background: | Synonyms: Protein FAM134C, FAM134C |
|---------------------|--|
| | Background: Chromosome 17 makes up over 2.5 % of the human genome with about 81 million |
| | bases encoding over 1,200 genes. Two key tumor suppressor genes are associated with |
| | chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for |
| | maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell |
| | death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li- |
| | Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, though specifically it is |
| | recognized as a genetic determinant of early onset breast cancer and predisposition to cancers |
| | of the ovary, colon, prostate gland and fallopian tubes. Chromosome 17 is also linked to |
| | neurofibromatosis, a condition characterized by neural and epidermal lesions, and dysregulated |
| | Schwann cell growth. Alexander disease, Birt-Hogg-Dube syndrome and Canavan disease are |
| | also associated with chromosome 17. The FAM134C gene product has been provisionally |
| | designated FAM134C pending further characterization. |
| Gene ID: | 162427 |
| UniProt: | Q86VR2 |
| Pathways: | SARS-CoV-2 Protein Interactome |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| | |

Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |