# antibodies -online.com





## anti-FAM160B1 antibody (AA 521-630) (Biotin)



#### Overview

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | FAM160B1  |
| Binding Specificity: | AA 521-630  |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This FAM160B1 antibody is conjugated to Biotin  |
| Application:         | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human FAM160B1 |
|-----------------------|--|
| Isotype:              | IgG  |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Chicken,Rabbit             |
| Purification:         | Purified by Protein A.                                       |

### **Target Details**

| Target:           | FAM160B1   |
|-------------------|--|
| Alternative Name: | FAM160B1 (FAM160B1 Products)   |
| Background:       | Synonyms: DKFZp686D10123, F16B1_HUMAN, Fam160b1, Family with sequence similarity 160 |

member B1, Hypothetical protein LOC57700, KIAA1600, Protein FAM160B1.

Background: FAM168A is a 244 amino acid protein that exists as three alternatively spliced isoforms and is encoded by a gene that maps to human chromosome 11, which makes up around 4% of human genomic DNA and is considered a gene and disease association dense chromosome. The chromosome 11 encoded Atm gene is important for regulation of cell cycle arrest and apoptosis following double strand DNA breaks. Atm mutation leads to the disorder known as ataxia-telangiectasia. The blood disorders Sickle cell anemia and thalassemia are caused by HBB gene mutations. Wilms' tumors, WAGR syndrome and Denys-Drash syndrome are associated with mutations of the WT1 gene. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are also associated with defects in chromosome 11.

#### **Application Details**

| Application Notes: | WB 1:300-5000 |
|--------------------|---------------|
|                    |               |

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

#### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C for 12 months.  |
| Expiry Date:       | 12 months  |