antibodies -online.com





anti-FAM21C antibody (AA 1201-1318) (Cy3)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | FAM21C |
| Binding Specificity: | AA 1201-1318 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FAM21C antibody is conjugated to Cy3 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human FAM21C |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Rat |
| Predicted Reactivity: | Human,Mouse,Cow,Pig,Chicken,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | FAM21C |
|-------------------|--------------------------|
| Alternative Name: | FAM21C (FAM21C Products) |

Target Details

| Background: | Synonyms: FAM21A, Family with sequence similarity 21 member C, Hypothetical protein | |
|---------------------|---|--|
| | LOC253725, KIAA0592, Protein FAM21C, FA21C_HUMAN. | |
| | Background: Playing a essential role in the fission of tubules that serve as transport | |
| | intermediates during endosome sorting, the WASH complex is present at the surface of | |
| | endosomes and functions to recruit and activate the Arp2/3 complex for induction of actin | |
| | polymerization. FAM21, Putative WASH complex subunit FAM21, is a component of the WASH | |
| | complex. Having undergone evolutionary duplication, four highly homologous family members | |
| | exist including FAM21A, FAM21B, FAM21C and FAM21D. FAM21 links the WASH complex to | |
| | endosomes and is required for WASM-dependent retromer-mediated sorting. Also, by directly | |
| | interacting with CapZ, FAM21 inhibits its anti-capping activity, thereby regulating actin | |
| | dynamics. | |
| Gene ID: | 253725 | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |
| | | |