

### Datasheet for ABIN896525

# anti-FAM3C antibody (AA 151-227) (Cy5)



| ()     | ve  | r\/i  |               | ۱۸/ |
|--------|-----|-------|---------------|-----|
| $\cup$ | V C | 1 / 1 | $\overline{}$ | ٧V  |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | FAM3C  |
| Binding Specificity: | AA 151-227   |
| Reactivity:          | Human, Rat   |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This FAM3C antibody is conjugated to Cy5   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |
| Product Details      |  |

#### Product Details

| Immunogen:            | KLH conjugated synthetic peptide derived from human FAM3C |  |
|-----------------------|---|--|
| Isotype:              | IgG   |  |
| Cross-Reactivity:     | Human, Rat  |  |
| Predicted Reactivity: | Mouse,Dog,Cow,Pig,Horse,Chicken                           |  |
| Purification:         | Purified by Protein A.                                    |  |

## Target Details

| Target:           | FAM3C                  |
|-------------------|------------------------|
| Alternative Name: | FAM3C (FAM3C Products) |

### **Target Details**

| Background:         | Synonyms: Fam3c, FAM3C_HUMAN, family with sequence similarity 3, member C, GS3786,ILEI,         |  |  |
|---------------------|---|--|--|
|                     | Interleukin-like EMT inducer, Predicted osteoblast protein, Protein FAM3C.                      |  |  |
|                     | Background: ILEI is a 227 amino acid, ubiquitously expressed protein containing an amino-       |  |  |
|                     | terminal signal peptide. Elevated levels of ILEI translation are observed in oncogenic, Ras-    |  |  |
|                     | transformed mammary epithelial cells and epithelial to mesenchymal transition (Emt) as well     |  |  |
|                     | as tumor growth and metastasis. Also, overexpression of ILEI results in loss of ZO-1, a protein |  |  |
|                     | involved in tight junctions, and expression of cytoplasmic E-cadherin, which has been shown to  |  |  |
|                     | influence loss of polarity and invasiveness. Due to this evidence, it is suspected that ILEI    |  |  |
|                     | cooperates with oncogenic Ras to cause TGF-independent Emt and its overexpression is            |  |  |
|                     | correlated with the invasion, metastasis and survival in a variety of cancers.                  |  |  |
| Gene ID:            | 10447   |  |  |
| Application Details |   |  |  |
| Application Notes:  | IF(IHC-P) 1:50-200  |  |  |
|                     | IF(IHC-F) 1:50-200  |  |  |
|                     | IF(ICC) 1:50-200  |  |  |
| Restrictions:       | For Research Use only   |  |  |
| Handling            |   |  |  |
| Format:             | Liquid  |  |  |
| Concentration:      | 1 μg/μL   |  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and     |  |  |
|                     | 50 % Glycerol.  |  |  |
| Preservative:       | ProClin   |  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be             |  |  |
|                     | handled by trained staff only.  |  |  |
| Storage:            | -20 °C  |  |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.               |  |  |
| Expiry Date:        | 12 months   |  |  |
|                     |   |  |  |