antibodies -online.com





anti-FAM59A antibody (AA 151-250) (Cy5)



Go to Product page

| \sim | | | | |
|--------|------|-----|-----|-----|
| | 11/6 | ٦r١ | /10 | ۱۸. |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | FAM59A |
| Binding Specificity: | AA 151-250 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FAM59A antibody is conjugated to Cy5 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human FAM59A | |
|-----------------------|--|--|
| Isotype: | IgG | |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Pig,Horse,Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | FAM59A | |
|--|--------------------------|--|
| Alternative Name: | FAM59A (FAM59A Products) | |
| Background: Synonyms: FA59A_HUMAN, fam59a, Family with sequence similarity 59, member A, GAF | | |

Gm944, Protein FAM59A.

Background: Encoding over 300 genes, chromosome 18 contains about 76 million bases. Trisomy 18, or Edwards syndrome, is the second most common trisomy after Downs syndrome. Symptoms of Edwards syndrome include low birth weight, a variety of physical development defects, heart deformations and breathing difficulty. Translocation between chromosome 18 and 14 is the most common translocation in cancers and occurs in follicular lymphomas. Niemann-Pick disease, hereditary hemorrhagic telangiectasia and erythropoietic protoporphyria are associated with chromosome 18. The TGF modulators, Smad2, Smad4 and Smad7 are encoded by chromosome 18. The FAM59A gene product has been provisionally designated FAM59A pending further characterization.

Gene ID: 64762

Pathways: EGFR Signaling Pathway

Application Details

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

| Format: | Liquid | |
|--------------------|--|--|
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |