

## Datasheet for ABIN896952

## anti-FCAR antibody (AA 131-230) (AbBy Fluor® 488)



Go to Product page

Overview

Quantity:	100 μL
Target:	FCAR
Binding Specificity:	AA 131-230
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FCAR antibody is conjugated to AbBy Fluor® 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human FCAR
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Purification:	Purified by Protein A.

## **Target Details**

Target:	FCAR
Alternative Name:	FCAR (FCAR Products)
Background:	Synonyms: CD89, Fc alpha receptor, FCAR, FCAR_HUMAN, IgA Fc receptor, Immunoglobulin

Expiry Date:

Target Details	
	alpha Fc receptor.  Background: Binds to the Fc region of immunoglobulins alpha. Mediates several functions including cytokine production. Tissue specificity, Isoform A.1, isoform A.2 and isoform A.3 are differentially expressed between blood and mucosal myeloid cells. Isoform A.1, isoform A.2 and isoform A.3 are expressed in monocytes. Isoform A.1 and isoform A.2 are expressed in alveolar macrophages, however only one isoform is expressed at alveolar macrophages surfaces.
Gene ID:	2204
Application Details	
Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

12 months