# antibodies -online.com





# anti-FDCSP antibody (AA 18-85) (Alexa Fluor 647)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	FDCSP (C4orf7)
Binding Specificity:	AA 18-85
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FDCSP antibody is conjugated to Alexa Fluor 647
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from rat FDC-SP
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Horse
Purification:	Purified by Protein A.

## Target Details

Target:	FDCSP (C4orf7)
Alternative Name:	Fdc-Sp (C4orf7 Products)

## **Target Details**

Background:	Synonyms: FDC, FDCs, Follicular dendritic cell, Follicular dendritic cell secreted protein,		
	Follicular dendritic cell, Follicular dendritic cell secreted peptide, FDC secreted protein.		
	Background: Follicular dendritic cells are cells that facilitate antigen recognition by B cells in		
	follicles. This kind of dendritic cell is not bone marrow derived and is not a kind of blood cell. It		
	is purely a resident of the follicles of secondary lymphoid organs. B cells form germinal centers		
	around follicular dendritic cells in lymphoid organs. Dendritic cells form from monocytes, white		
	blood cells which circulate in the body and, depending on the right signal, can turn into dendritic		
	cells or macrophages. The monocytes in turn are formed from stem cells in the bone marrow. In		
	normal tissue this antibody identifies dendritic cells and a proportion of B lymphocytes.		
	Evidence in pathological tissue and functional studies suggests it binds to an epitope		
	expressed by antigen presenting cells. Within the macrophage/dendritic cell populations the		
	epitope seen by RFD1 is coexpressed with RFD7 by a subset of cells that exhibit suppressive		
	activity on T cell stimulation.		
Gene ID:	494224		
UniProt:	Q5R1M6		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		

		1.	
$\vdash$	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months