

## Datasheet for ABIN897190

## anti-FGG antibody (AA 151-250) (Cy5)



Go to Product page

| ()      | ve  | r\/i | Δ | ۱۸/ |
|---------|-----|------|---|-----|
| $\circ$ | V C | 1 V  |   | v v |

Alternative Name:

| Quantity:             | 100 μL   |  |
|-----------------------|--|--|
| Target:               | FGG  |  |
| Binding Specificity:  | AA 151-250   |  |
| Reactivity:           | Human, Mouse, Rat  |  |
| Host:                 | Rabbit   |  |
| Clonality:            | Polyclonal   |  |
| Conjugate:            | This FGG antibody is conjugated to Cy5   |  |
| Application:          | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |
| Product Details       |  |  |
| Immunogen:            | KLH conjugated synthetic peptide derived from human Fibrinogen gamma chain   |  |
| Isotype:              | IgG  |  |
| Cross-Reactivity:     | Human, Mouse, Rat  |  |
| Predicted Reactivity: | Dog,Horse,Rabbit   |  |
| Purification:         | Purified by Protein A.   |  |
| Target Details        |  |  |
| Target:               | FGG  |  |
|                       |  |  |

Fibrinogen gamma chain (FGG Products)

## **Target Details**

| Background:         | Synonyms: Fibrinogen gamma chain, FGG, PRO2061   |  |
|---------------------|--|--|
|                     | Background: Together with fibrinogen alpha (FGA) and fibrinogen beta (FGB), polymerizes to           |  |
|                     | form an insoluble fibrin matrix. Has a major function in hemostasis as one of the primary            |  |
|                     | components of blood clots. In addition, functions during the early stages of wound repair to         |  |
|                     | stabilize the lesion and guide cell migration during re-epithelialization. Was originally thought to |  |
|                     | be essential for platelet aggregation, based on in vitro studies using anticoagulated blood.         |  |
|                     | However, subsequent studies have shown that it is not absolutely required for thrombus               |  |
|                     | formation in vivo. Enhances expression of SELP in activated platelets via an ITGB3-dependent         |  |
|                     | pathway. Maternal fibrinogen is essential for successful pregnancy. Fibrin deposition is also        |  |
|                     | associated with infection, where it protects against IFNG-mediated hemorrhage. May also              |  |
|                     | facilitate the antibacterial immune response via both innate and T-cell mediated pathways.           |  |
| Gene ID:            | 2266   |  |
| UniProt:            | P02679   |  |
| Application Details |  |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |  |
|                     | IF(IHC-F) 1:50-200   |  |
|                     | IF(ICC) 1:50-200   |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Concentration:      | 1 μg/μL  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and          |  |
|                     | 50 % Glycerol.   |  |
| Preservative:       | ProClin  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be                  |  |
|                     | handled by trained staff only.   |  |
| Storage:            | -20 °C   |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                    |  |
| Expiry Date:        | 12 months  |  |
|                     |  |  |