antibodies -online.com





anti-FRMD8 antibody (AA 65-170) (HRP)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	FRMD8	
Binding Specificity:	AA 65-170	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This FRMD8 antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FRMD8
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig
Purification:	Purified by Protein A.

Target Details

Target:	FRMD8
Alternative Name:	FRMD8 (FRMD8 Products)

Target Details

Background:	Synonyms: FERM domain containing protein 8, FKSG44, FRMD8_HUMAN.
	Background: FERM domains are roughly 150 amino acids in length and are found in a number
	of cytoskeletal-associated proteins such as Ezrin, Radixin, Moesin and 4.1 (erythrocyte
	membrane protein band 4.1), where they provide a link between cytoskeletal signals and
	membrane dynamics. FRMD8 (FERM domain-containing protein 8), also known as FKSG44, is a
	464 amino acid protein containing one FERM domain. Existing as two alternatively spliced
	isoforms, the gene encoding FRMD8 maps to human chromosome 11q13.1. With
	approximately 135 million base pairs and 1,400 genes, chromosome 11 makes up around 4 $\%$
	of human genomic DNA and is considered a gene and disease association dense chromosome.
	Wilms' tumors, WAGR syndrome and Denys-Drash syndrome are associated with mutations of
	the WT1 gene. Jervell and Lange-Nielsen syndrome, Jacobsen syndrome, Niemann-Pick
	disease, hereditary angioedema and Smith-Lemli-Opitz syndrome are also associated with
	defects in chromosome 11.
Gene ID:	83786
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	 1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish
	peroxidase.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months