

Datasheet for ABIN898036

**anti-GBL antibody (AA 181-290) (AbBy Fluor® 647)**[Go to Product page](#)

## Overview

|                      |                                                                                                         |
|----------------------|---------------------------------------------------------------------------------------------------------|
| Quantity:            | 100 µL                                                                                                  |
| Target:              | GBL                                                                                                     |
| Binding Specificity: | AA 181-290                                                                                              |
| Reactivity:          | Mouse, Rat                                                                                              |
| Host:                | Rabbit                                                                                                  |
| Clonality:           | Polyclonal                                                                                              |
| Conjugate:           | This GBL antibody is conjugated to AbBy Fluor® 647                                                      |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## Product Details

|                       |                                                                                 |
|-----------------------|---------------------------------------------------------------------------------|
| Immunogen:            | KLH conjugated synthetic peptide derived from human G protein beta subunit like |
| Isotype:              | IgG                                                                             |
| Cross-Reactivity:     | Mouse, Rat                                                                      |
| Predicted Reactivity: | Human,Dog,Cow,Sheep,Pig                                                         |
| Purification:         | Purified by Protein A.                                                          |

## Target Details

|                   |                                                              |
|-------------------|--------------------------------------------------------------|
| Target:           | GBL                                                          |
| Alternative Name: | G protein beta subunit like ( <a href="#">GBL Products</a> ) |

## Target Details

|             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
|-------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Background: | <p>Synonyms: G protein beta subunit like, G protein beta subunit-like, Gable, GBL, GBL protein, LST8_HUMAN, Mammalian lethal with SEC13 protein 8, MGC111011, mLST8, Protein GbetaL, Target of rapamycin complex subunit LST8, TORC subunit LST8.</p> <p>Background: Subunit of both mTORC1 and mTORC2, which regulates cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids. Growth factor-stimulated mTORC1 activation involves a AKT1-mediated phosphorylation of TSC1-TSC2, which leads to the activation of the RHEB GTPase that potentially activates the protein kinase activity of mTORC1. Amino-acid-signaling to mTORC1 requires its relocalization to the lysosomes mediated by the Ragulator complex and the Rag GTPases. Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. mTORC1 phosphorylates EIF4EBP1 and releases it from inhibiting the elongation initiation factor 4E (eIF4E). mTORC1 phosphorylates and activates S6K1 at 'Thr-389', which then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation. Tissue specificity: Broadly expressed, with highest levels in skeletal muscle, heart and kidney.</p> |
|-------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

|           |                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Pathways: | <a href="#">PI3K-Akt Signaling</a> , <a href="#">RTK Signaling</a> , <a href="#">Fc-epsilon Receptor Signaling Pathway</a> , <a href="#">EGFR Signaling Pathway</a> , <a href="#">Neurotrophin Signaling Pathway</a> , <a href="#">Regulation of Actin Filament Polymerization</a> , <a href="#">Autophagy</a> , <a href="#">CXCR4-mediated Signaling Events</a> , <a href="#">BCR Signaling</a> , <a href="#">Warburg Effect</a> |
|-----------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

## Application Details

|                    |                                                                             |
|--------------------|-----------------------------------------------------------------------------|
| Application Notes: | <p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p> |
|--------------------|-----------------------------------------------------------------------------|

|               |                       |
|---------------|-----------------------|
| Restrictions: | For Research Use only |
|---------------|-----------------------|

## Handling

|                    |                                                                                                                    |
|--------------------|--------------------------------------------------------------------------------------------------------------------|
| Format:            | Liquid                                                                                                             |
| Concentration:     | 1 µg/µL                                                                                                            |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.        |
| Preservative:      | ProClin                                                                                                            |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |

Handling

---

|                  |                                                                                   |
|------------------|-----------------------------------------------------------------------------------|
| Storage:         | -20 °C                                                                            |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date:     | 12 months                                                                         |