

Datasheet for ABIN898894

anti-GIMAP6 antibody (AA 25-130) (HRP)



_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Overview			
Quantity:	100 μL		
Target:	GIMAP6		
Binding Specificity:	AA 25-130		
Reactivity:	Mouse		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This GIMAP6 antibody is conjugated to HRP		
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)),		
	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))		
Product Details			
Immunogen:	KLH conjugated synthetic peptide derived from mouse GIMAP6		
Isotype:	IgG		
Cross-Reactivity:	Mouse		
Predicted Reactivity:	Rat		
Purification:	Purified by Protein A.		
Target Details			
Target:	GIMAP6		
Alternative Name:	GIMAP6 (GIMAP6 Products)		

Target Details

immune associated nucleotide 2, IAN-2, IAN-6, IAN6, Immune associated nucleotide associated nucleotide 6, Immunity associated nucleotide 6 protein, Immunity-associated nucleotide 6 protein. Immunity-associated nucleotide 6 protein, Immunity-associated nucleotide 6 protein. Immunity-associated nucleotide 6 protein. Background: Highly expressed in splean, lyngh packground and placenta. Expression in tissues. Immunity-associated nucleotide 6 protein. Background: Highly expressed in splean, lyngh phackground and placenta. Expressed in splean and pl	ranger Betane		
Application Details Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, w handled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of hoperoxidase. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Background:	Background: Highly expressed in spleen, lymph nodes, lung and placenta. Expressed at moderate level in thymus, kidney, heart and digestive tract. Weakly expressed in other lymphoid	
Application Notes: WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, w handled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of hoperoxidase. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Gene ID:	474344	
IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, whandled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of hoperoxidase. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Application Details		
Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, w handled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of he peroxidase. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Application Notes:	IHC-P 1:200-400	
Format: Liquid Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, whandled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of hoperoxidase. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Restrictions:	For Research Use only	
Concentration: 1 μg/μL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, w handled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of hoperoxidase. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Handling		
Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % 50 % Glycerol. Preservative: ProClin This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, whandled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of hoperoxidase. Storage: -20 °C Storage Comment: Store at -20 °C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Format:	Liquid	
Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, we handled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of hoperoxidase. Storage: -20 °C Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Concentration:	1 μg/μL	
Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, whandled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of hoperoxidase. Storage: -20 °C Storage Comment: Store at -20 °C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
handled by trained staff only. Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of hoperoxidase. Storage: -20 °C Storage Comment: Store at -20 °C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Preservative:	ProClin	
peroxidase. Storage: -20 °C Storage Comment: Store at -20 °C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
	Storage:	-20 °C	
Expiry Date: 12 months	Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
	Expiry Date:	12 months	