# antibodies -online.com





## anti-KCNJ3 antibody (AA 81-180) (Alexa Fluor 350)



Go to Product page

| ( ) | 1/0 | r\ /1 | 014 |   |
|-----|-----|-------|-----|---|
| ( ) | ve  | I V I | -v  | V |
|     |     |       |     |   |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | KCNJ3  |
| Binding Specificity: | AA 81-180  |
| Reactivity:          | Mouse, Rat   |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This KCNJ3 antibody is conjugated to Alexa Fluor 350   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human GIRK1 |
|-----------------------|---|
| Isotype:              | IgG   |
| Cross-Reactivity:     | Mouse, Rat  |
| Predicted Reactivity: | Human,Dog,Cow,Pig,Rabbit,Guinea Pig                       |
| Purification:         | Purified by Protein A.                                    |

#### **Target Details**

| Target:           | KCNJ3                  |
|-------------------|------------------------|
| Alternative Name: | GIRK1 (KCNJ3 Products) |

### **Target Details**

| Background:         | Synonyms: KGA, GIRK1, KIR3.1, G protein-activated inward rectifier potassium channel 1, GIRK-  |  |  |
|---------------------|--|--|--|
|                     | 1, Inward rectifier K(+) channel Kir3.1, Potassium channel, inwardly rectifying subfamily J  |  |  |
|                     | member 3, KCNJ3  |  |  |
|                     | Background: This potassium channel is controlled by G proteins. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium, as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward |  |  |
|                     |  |  |  |
|                     |  |  |  |
|                     |  |  |  |
|                     |  |  |  |
| Gene ID:            | 3760   |  |  |
| UniProt:            | P48549   |  |  |
| A 1: 1: D 1:1       |  |  |  |
| Application Details |  |  |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |  |  |
|                     | IF(IHC-F) 1:50-200   |  |  |
|                     | IF(ICC) 1:50-200   |  |  |
| Restrictions:       | For Research Use only  |  |  |
| Handling            |  |  |  |
| Format:             | Liquid   |  |  |
| Concentration:      | 1 μg/μL  |  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and  |  |  |
|                     | 50 % Glycerol.   |  |  |
| Preservative:       | ProClin  |  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be  |  |  |
|                     | handled by trained staff only.   |  |  |
| Storage:            | -20 °C   |  |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.  |  |  |
| Expiry Date:        | 12 months  |  |  |
|                     |  |  |  |