antibodies -online.com





anti-GPRIN2 antibody (AA 251-350) (HRP)



Go to Product page

| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | GPRIN2 |
| Binding Specificity: | AA 251-350 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GPRIN2 antibody is conjugated to HRP |
| Application: | ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human GPRIN2 |
|-----------------------|--|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Pig,Horse |
| Purification: | Purified by Protein A. |

Target Details

| Target: | GPRIN2 | |
|-------------------|--|--|
| Alternative Name: | GPRIN2 (GPRIN2 Products) | |
| Background: | Synonyms: G protein-regulated inducer of neurite outgrowth 2, Gm286, G protein-regulated | |

inducer of neurite outgrowth 2, GPRIN2, GRIN2, GRIN2_HUMAN, MGC15171, mKIAA0514, RGD1308922.

Background: G protein-coupled receptors (GPCRs) represent a large superfamily of cell-surface receptors that are involved in a multitude of physiological processes such as perception of sensory information, modulation of synaptic transmission, hormone release/actions, regulation of cell contraction/migration and cell growth/differentiation. GPCRs interact with G proteins (heterotrimeric GTPases) to synthesize intracellular second messengers, such as diacylglycerol, cyclic AMP, inositol phosphates and calcium ions. Their diverse biological functions range from vision and olfaction to neuronal and endocrine signaling and are involved in many pathological conditions. GRIN2 (G protein-regulated inducer of neurite outgrowth 2), also known as GPRIN2, is a 458 amino acid protein that is expressed in cerebellum and is thought to play a role in neurite outgrowth. GRIN2 interacts with activated G?oand G?, and is encoded by a gene that maps to human chromosome 10q11.22.

Gene ID:

9721

Application Details

| Application Notes: | IHC-P 1:200-400 |
|--------------------|-----------------|
| | |

IHC-F 1:100-500

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Handling Advice: | Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |

| 1 1 | | |
|----------------|----------|-------|
| \blacksquare | α | lina |
| | land | 1111(|
| | I G I G | 11119 |
| | | |

Expiry Date:

12 months