

[Go to Product page](#)

Datasheet for ABIN900188  
**anti-HBSAg antibody (Alexa Fluor 488)**

## Overview

Quantity:	100 µL
Target:	HBSAg (HBsAg)
Reactivity:	Virus, Human, Mouse
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This HBSAg antibody is conjugated to Alexa Fluor 488
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

## Product Details

Immunogen:	Human Hepatitis B Surface Antigen
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Virus
Predicted Reactivity:	Bee
Purification:	Purified by Protein A.

## Target Details

Target:	HBSAg (HBsAg)
Alternative Name:	Hbsag ( <a href="#">HBSAg Products</a> )
Target Type:	Viral Protein

## Target Details

Background:	<p>Synonyms: HBsAg, HBV major surface antigen, HBV surface antigen, Hepatitis B Virus major surface antigen, Major surface antigen, S, Human Hepatitis B Surface Antigen</p> <p>Background: Hepatitis B Virus (HBV) infection induces a disease state characterised by liver damage, inflammation and viral persistence. Infection also increases the risk of hepatocellular carcinoma. HBV belongs to the Hepadnaviridae family of viruses. Its genome consists of partially double stranded circular DNA. The DNA is enclosed in a nucleocapsid, or core antigen (HBcAg), which is surrounded by a spherical envelope (surface antigen or HBsAg). The core antigen shares its sequences with the e antigen (HBeAg) but no cross reactivity between the two proteins has been observed. The HBV genome also encodes a DNA polymerase that also acts as a reverse transcriptase. Hepatitis B infection is normally diagnosed from serological tests that detect HBsAg but as the disease progresses this antigen may no longer be present in the blood and tests for HBcAg are used. If HBsAg can be detected in the blood for longer than six months, chronic hepatitis B is diagnosed. The antigenic determinant of the protein moiety of the HBsAg determines specific characteristics of different serotypes and provides the basis of immunodetection. HBsAg has antigenic heterogeneity, specifically, two pairs of sub specific determinants, d/y and w/r allow the following combinations: adw, ayw, adr, ayr.</p>
-------------	--

UniProt: [Q67926](#)

## Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

## Handling

---

Storage:	-20 °C
----------	--------

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
------------------	---

Expiry Date:	12 months
--------------	-----------