



[Go to Product page](#)

Datasheet for ABIN900793

## anti-HIRA antibody (AA 251-350) (AbBy Fluor® 350)

### Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | HIRA   |
| Binding Specificity: | AA 251-350   |
| Reactivity:          | Mouse  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This HIRA antibody is conjugated to AbBy Fluor® 350  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

### Product Details

|                       |  |
|-----------------------|--|
| Immunogen:            | KLH conjugated synthetic peptide derived from human HIRA/DGGR1 |
| Isotype:              | IgG  |
| Cross-Reactivity:     | Mouse  |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Pig,Horse,Rabbit                             |
| Purification:         | Purified by Protein A.   |

### Target Details

|                   |  |
|-------------------|--|
| Target:           | HIRA                                   |
| Alternative Name: | HIRA ( <a href="#">HIRA Products</a> ) |

## Target Details

---

**Background:** Synonyms: DGCR1, DGGR 1, DGGR1, DiGeorge critical region gene 1, HIR, HIR histone cell cycle regulation defective homolog A, HIRA, HIRA protein, HIRA\_HUMAN, Protein HIRA, TUP 1, TUP1, TUP1 like enhancer of split protein 1, TUP1-like enhancer of split protein 1, TUPLE 1, TUPLE1.

Background: The HIRA gene encodes a histone chaperone that preferentially places the variant histone H3.3 in nucleosomes. Orthologs of this gene in flies, yeast and plants are necessary for the formation of transcriptionally silent heterochromatin. This gene plays an important role in the formation of the senescence-associated heterochromatin foci. These foci likely mediate the irreversible cell cycle changes that occur in senescent cells. It is considered the primary candidate gene in some haploinsufficiency syndromes such as DiGeorge syndrome, and insufficient production of the gene may disrupt normal embryonic development.

---

**Gene ID:** 7290

---

**Pathways:** [Chromatin Binding](#)

## Application Details

---

**Application Notes:** IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

---

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

---

**Concentration:** 1 µg/µL

---

**Buffer:** Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

**Preservative:** ProClin

---

**Precaution of Use:** This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

**Storage:** -20 °C

---

**Storage Comment:** Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

**Expiry Date:** 12 months