antibodies .- online.com







anti-HMGCL antibody (AA 41-140) (AbBy Fluor® 555)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	HMGCL
Binding Specificity:	AA 41-140
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMGCL antibody is conjugated to AbBy Fluor® 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HMGCL
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	HMGCL
Alternative Name:	HMGCL (HMGCL Products)

Target Details

Background:	Synonyms: 3 hydroxy 3 methylglutaryl CoA lyase, 3 hydroxy 3 methylglutaryl Coenzyme A lyase,	
	3 hydroxymethyl 3 methylglutaryl Coenzyme A lyase hydroxymethylglutaricaciduria, 3	
	hydroxymethyl 3 methylglutaryl Coenzyme A lyase, 3-hydroxy-3-methylglutarate-CoA lyase, HL,	
	HMG CoA lyase, HMG CoA Lyase Deficiency, HMG-CoA lyase, HMGCL, HMGCL_HUMAN,	
	Hydroxymethylglutaricaciduria, Hydroxymethylglutaryl CoA lyase, Hydroxymethylglutaryl CoA	
	lyase mitochondrial, Hydroxymethylglutaryl-CoA lyase, mitochondrial, MS725,	
	OTTHUMP0000044830.	
	Background: Hydroxymethylglutaryl-CoA lyase (HMGCL) is found in fibroblasts, liver and	
	lymphoblasts. It has a role in ketogenesis and leucine catabolism. Defects in HMGCL are the	
	cause of 3-hydroxy-3-methylglutaryl-CoA lyase deficiency (hydroxymethylglutaricaciduria), an	
	autosomal recessive disease which can lead to hypoglycemia and coma.	
Gene ID:	2835	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	