## antibodies -online.com





## anti-HRPT2 antibody (AA 61-160) (Alexa Fluor 350)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	HRPT2 (CDC73)
Binding Specificity:	AA 61-160
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HRPT2 antibody is conjugated to Alexa Fluor 350
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human HRPT2/Parafibromin
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig
Purification:	Purified by Protein A.

## **Target Details**

Target:	HRPT2 (CDC73)
Alternative Name:	Parafibromin (CDC73 Products)
Background:	Synonyms: C1orf28,CDC 73,CDC73,Cell division cycle 73,Cell division cycle 73 Paf1/RNA

Expiry Date:

Target Details	
	polymerase II complex component homolog, Cell division cycle protein 73 homolog, FLJ23316, HPT-JT, HRPT 2, Hyperparathyroidism 2 with jaw tumor, Hyperparathyroidism 2, Hyperparathyroidism 2 protein, Parafibromin.  Background: The HRPT2 gene product, parafibromin, is a tumor suppressor protein that is part of the human Paf1 complex. The yeast counterpart to this complex is part of the RNA polymerase II complex as well, and is important for histone modification and connections to posttranscriptional events. Human parafibromin also associates with the RNA polymerase II large subunit.
Gene ID:	79577
Pathways:	Cellular Response to Molecule of Bacterial Origin, Stem Cell Maintenance
Application Details	
Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

12 months