antibodies .- online.com





anti-HSD17B2 antibody (AA 151-250) (Alexa Fluor 350)



Overview

Quantity:	100 μL
Target:	HSD17B2
Binding Specificity:	AA 151-250
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HSD17B2 antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HSD17B2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat
Purification:	Purified by Protein A.

Target Details

Target:	HSD17B2
Alternative Name:	HSD17B2 (HSD17B2 Products)

Target Details

-	
Background:	Synonyms: HSD17, SDR9C2, EDH17B2, Estradiol 17-beta-dehydrogenase 2, 17-beta-
	hydroxysteroid dehydrogenase type 2, 17-beta-HSD 2, 20 alpha-hydroxysteroid dehydrogenase,
	20-alpha-HSD, E2DH, Microsomal 17-beta-hydroxysteroid dehydrogenase, Short chain
	dehydrogenase/reductase family 9C member 2, Testosterone 17-beta-dehydrogenase,
	HSD17B2
	Background: Capable of catalyzing the interconversion of testosterone and androstenedione, as
	well as estradiol and estrone. Also has 20-alpha-HSD activity. Uses NADH while EDH17B3 uses
	NADPH.
Gene ID:	3294
UniProt:	P37059
Pathways:	Steroid Hormone Biosynthesis
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months