antibodies - online.com







anti-ILP-2 antibody (AA 141-236) (Cy5)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	ILP-2 (BIRC8)
Binding Specificity:	AA 141-236
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ILP-2 antibody is conjugated to Cy5
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ILP2	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human, Mouse, Chicken, Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	ILP-2 (BIRC8)
Alternative Name:	ILP2 (BIRC8 Products)

Target Details

Background:

Synonyms: Baculoviral IAP repeat containing 8, Baculoviral IAP repeat containing protein 8, Baculoviral IAP repeat-containing protein 8, BIRC 8, BIRC8, BIRC8 protein, BIRC8_HUMAN, hILP 2, hILP2, IAP like protein 2, IAP-like protein 2, ILP 2, ILP-2, Inhibitor of apoptosis like protein 2, Inhibitor of apoptosis-like protein 2, Testis specic inhibitor of apoptosis, Testis-specic inhibitor of apoptosis.

Background: Protects against apoptosis mediated by BAX.Apoptosis, or programmed cell death, is related to many diseases, such as cancer. Apoptosis is triggered by a variety of stimuli including members in the TNF family and prevented by the inhibitor of apoptosis (IAP) proteins. IAP proteins form a conserved gene family including IAP, XIAP/ILP-1/MIHA, and Livin/KIAP that bind to and inhibits specific proteases. A novel member in the IAP protein family was recently identified and designated ILP-2 for IAP-like protein-2 (1). ILP-2 has high homology to ILP-1, but is encoded by a distinct gene that is solely expressed in testis of tested normal human tissues (1). ILP-2, unlike ILP-1, has no inhibitory effect on Fas and TNF induced apoptosis, but potently inhibits apoptosis induced by overexpression of Bax or by coexpression of caspase-9 with Apaf-1. ILP-2 interacts with the processed caspase-9. These results suggest that ILP-2 is a novel IAP family member with restricted specificity for caspase-9.

Gene ID:

112401

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months