

[Go to Product page](#)

Datasheet for ABIN902692

anti-Importin 9 antibody (AA 601-729) (Biotin)

Overview

Quantity:	100 µL
Target:	Importin 9 (IPO9)
Binding Specificity:	AA 601-729
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Importin 9 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Importin 9
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	Importin 9 (IPO9)
Alternative Name:	Importin 9 (IPO9 Products)

Target Details

Background:	<p>Synonyms: B cell antigen receptor Ig beta associated protein 1, BPM 90, BPM L, BPM-L, BPM90, BPML, IBAP 1, Imp 9, Importin 9, Importin9, Importin-9, Novel centrosomal protein RanBPM, RAN binding protein 9, Ran binding protein centrosomal, Ran binding protein M, Ran BP9, Ran-binding protein 9, Ran-binding protein M, RANB9_HUMAN, RanBP 7, RANBP 9, RanBP7, RanBP9, RanBPM.</p> <p>Background: May act as an adapter protein to couple membrane receptors to intracellular signaling pathways. May be involved in signaling of ITGB2/LFA-1 and other integrins. Enhances HGF-MET signaling by recruiting Sos and activating the Ras pathway. Involved in activation of androgen and glucocorticoid receptor in the presence of their cognate hormones. Stabilizes TP73 isoform Alpha, probably by inhibiting its ubiquitination, and increases its proapoptotic activity. Inhibits the kinase activity of DYRK1A and DYRK1B. Inhibits FMR1 binding to RNA. Tissue specificity: Ubiquitously expressed, with highest levels in testes, placenta, heart, and muscle, and lowest levels in lung. Within the brain, expressed predominantly by neurons in the gray matter of cortex, the granular layer of cerebellum and the Purkinje cells.</p>
-------------	--

Gene ID:	10048
Pathways:	Protein targeting to Nucleus

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C for 12 months.
------------------	-------------------------------

Expiry Date:	12 months
--------------	-----------