antibodies - online.com







anti-IRGM antibody (AA 51-150) (AbBy Fluor® 350)



| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | IRGM | |
| Binding Specificity: | AA 51-150 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This IRGM antibody is conjugated to AbBy Fluor® 350 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human IRGM | |
|-----------------------|--|--|
| Isotype: IgG | | |
| Predicted Reactivity: | activity: Human,Mouse,Rat,Dog | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | IRGM | |
|-------------------|---|--|
| Alternative Name: | IRGM (IRGM Products) | |
| Background: | Synonyms: Irgm, I1, Iigp3, Immunity related GTPase family M protein 1, Immunity related | |

GTPase family, M, Interferon inducible protein 1, IRGM1, LRG 47, LRG47, IRGM_HUMAN.

Background: IRGM belongs to a family of interferon-gamma inducible GTPases, belonging to the GTPase superfamily, and are selectively induced by IFN-gamma or bacterial lipopolysaccharide (LPS) stimulation. IRGM is primarily expressed in all cells derived from B-cell lineages, and is highly expressed in macrophages following IFN-gamma stimulation. IRGM has been shown to be required for host defences against a broad range of intracellular pathogens. Specifically, IRGM deficient mice show defects with intracellular microbial killing, phagosome maturation and autophagy. More recently IRGM has been shown to inhibit baseline hematopoietic proliferation and is required for a normal hematopoietic stem cell response to chemical and infectious stimuli.

Gene ID:

345611

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 | |
|--------------------|--------------------|--|
| | IF(IHC-F) 1:50-200 | |

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |