

Datasheet for ABIN903244

**anti-KDM3B antibody (AA 1601-1761) (AbBy Fluor® 488)**[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	KDM3B
Binding Specificity:	AA 1601-1761
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KDM3B antibody is conjugated to AbBy Fluor® 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human JMJD1B
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	KDM3B
Alternative Name:	JMJD1B ( <a href="#">KDM3B Products</a> )

## Target Details

---

Background:	<p>Synonyms: JHDM2B, JmjC domain containing histone demethylation protein 2B, JmjC domain-containing histone demethylation protein 2B , jmjd1b, Jumonji domain containing protein 1B, JUMONJI DOMAIN-CONTAINING PROTEIN 1B , KDM3B, KDM3B_HUMAN, Lysine-specific demethylase 3B, Nuclear protein 5qNCA.</p> <p>Background: JMJD1B (jumonji domain containing 1B), also known as KDM3B, 5qNCA (5q Nuclear Co-Activator) or C5orf7, is a member of the JHDM2 histone demethylase family of proteins. Expressed in a wide variety of tissues, JMJD1B localizes to the nucleus and contains one JMJC domain and a C-terminal zinc finger motif. JMJD1B functions as a histone demethylase and, using iron as a cofactor, demethylates lysine-9 of Histone H3. This suggests that JMJD1B plays a central role in the histone code. The gene encoding human JMJD1B is located within the 5q region of the genome that is often deleted in myeloid leukemias and myelodysplasias. This implies that JMJD1B may function as a tumor suppressor of myeloid leukemia. Ectopic expression of JMJD1B exhibits growth suppressive activities, further supporting a role for JMJD1B in tumor suppression.</p>
-------------	---

---

Gene ID:	51780
----------	-------

---

Pathways:	<a href="#">Warburg Effect</a>
-----------	--------------------------------

## Application Details

---

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
--------------------	--

---

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

---

Format:	Liquid
---------	--------

---

Concentration:	1 µg/µL
----------------	---------

---

Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
---------	---

---

Preservative:	ProClin
---------------	---------

---

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
--------------------	--

---

Storage:	-20 °C
----------	--------

---

## Handling

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months