antibodies .- online.com







anti-KCNMB1 antibody (AA 10-80) (Cy5)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	KCNMB1	
Binding Specificity:	AA 10-80	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This KCNMB1 antibody is conjugated to Cy5	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human KCNMB1	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	

Target Details

Target:	KCNMB1	
Alternative Name:	KCNMB1 (KCNMB1 Products)	

Target Details

Precaution of Use:

Storage Comment:

Storage:

Target Details		
Background:	Synonyms: Calcium-activated potassium channel subunit beta-1, KCNMB1, BK channel subunit	
	beta-1, Charybdotoxin receptor subunit beta-1, K(VCA)beta-1, Maxi K channel subunit beta-1,	
	Slo-beta-1, Slo-beta	
	Background: Regulatory subunit of the calcium activated potassium KCNMA1 (maxiK) channel.	
	Modulates the calcium sensitivity and gating kinetics of KCNMA1, thereby contributing to	
	KCNMA1 channel diversity. Increases the apparent Ca2+/voltage sensitivity of the KCNMA1	
	channel. It also modifies KCNMA1 channel kinetics and alters its pharmacological properties. It	
	slows down the activation and the deactivation kinetics of the channel. Acts as a negative	
	regulator of smooth muscle contraction by enhancing the calcium sensitivity to KCNMA1. Its	
	presence is also a requirement for internal binding of the KCNMA1 channel opener	
	dehydrosoyasaponin I (DHS-1) triterpene glycoside and for external binding of the agonist	
	hormone 17-beta-estradiol (E2). Increases the binding activity of charybdotoxin (CTX) toxin to	
	KCNMA1 peptide blocker by increasing the CTX association rate and decreasing the	
	dissociation rate.	
Gene ID:	3779	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

handled by trained staff only.

-20 °C

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Н	and	linc

Expiry Date:

12 months