# antibodies .- online.com







## anti-KCNMB1 antibody (AA 10-80) (Cy5.5)



$\sim$				
	11/6	٦r١	/10	۱۸.

Quantity:	100 μL
Target:	KCNMB1
Binding Specificity:	AA 10-80
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCNMB1 antibody is conjugated to Cy5.5
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human KCNMB1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

### **Target Details**

Target:	KCNMB1
Alternative Name:	KCNMB1 (KCNMB1 Products)

#### **Target Details**

Larget Details	
Background:	Synonyms: Calcium-activated potassium channel subunit beta-1, KCNMB1, BK channel subunit beta-1, Charybdotoxin receptor subunit beta-1, K(VCA)beta-1, Maxi K channel subunit beta-1, Slo-beta-1, Slo-beta  Background: Regulatory subunit of the calcium activated potassium KCNMA1 (maxiK) channel. Modulates the calcium sensitivity and gating kinetics of KCNMA1, thereby contributing to KCNMA1 channel diversity. Increases the apparent Ca2+/voltage sensitivity of the KCNMA1 channel. It also modifies KCNMA1 channel kinetics and alters its pharmacological properties. It slows down the activation and the deactivation kinetics of the channel. Acts as a negative regulator of smooth muscle contraction by enhancing the calcium sensitivity to KCNMA1. Its presence is also a requirement for internal binding of the KCNMA1 channel opener dehydrosoyasaponin I (DHS-1) triterpene glycoside and for external binding of the agonist
	hormone 17-beta-estradiol (E2). Increases the binding activity of charybdotoxin (CTX) toxin to
	KCNMA1 peptide blocker by increasing the CTX association rate and decreasing the
	dissociation rate.
Gene ID:	3779
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

-20 °C

Storage:

Storage Comment:

		1.	
$\vdash$	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months