

[Go to Product page](#)

Datasheet for ABIN903524

**anti-KCNMB1 antibody (AA 10-80) (Biotin)**

## Overview

Quantity:	100 µL
Target:	KCNMB1
Binding Specificity:	AA 10-80
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCNMB1 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human KCNMB1
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse, Cow, Sheep, Pig, Horse
Purification:	Purified by Protein A.

## Target Details

Target:	KCNMB1
Alternative Name:	KCNMB1 ( <a href="#">KCNMB1 Products</a> )

## Target Details

Background:	<p>Synonyms: Calcium-activated potassium channel subunit beta-1, KCNMB1, BK channel subunit beta-1, Charybdotoxin receptor subunit beta-1, K(VCA)beta-1, Maxi K channel subunit beta-1, Slo-beta-1, Slo-beta</p> <p>Background: Regulatory subunit of the calcium activated potassium KCNMA1 (maxiK) channel. Modulates the calcium sensitivity and gating kinetics of KCNMA1, thereby contributing to KCNMA1 channel diversity. Increases the apparent Ca<sup>2+</sup>/voltage sensitivity of the KCNMA1 channel. It also modifies KCNMA1 channel kinetics and alters its pharmacological properties. It slows down the activation and the deactivation kinetics of the channel. Acts as a negative regulator of smooth muscle contraction by enhancing the calcium sensitivity to KCNMA1. Its presence is also a requirement for internal binding of the KCNMA1 channel opener dehydrosoyasaponin I (DHS-1) triterpene glycoside and for external binding of the agonist hormone 17-beta-estradiol (E2). Increases the binding activity of charybdotoxin (CTX) toxin to KCNMA1 peptide blocker by increasing the CTX association rate and decreasing the dissociation rate.</p>
-------------	---

Gene ID:	3779
----------	------

## Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months