# antibodies -online.com







# anti-ARHGEF12 antibody (AA 515-615) (Alexa Fluor 350)



| ( ) | 11/0   | K\ /   | iew         |
|-----|--------|--------|-------------|
|     | $\cup$ | 'I V/I | $I \cap VV$ |
|     |        |        |             |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | ARHGEF12   |  |
| Binding Specificity: | AA 515-615   |  |
| Reactivity:          | Mouse  |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This ARHGEF12 antibody is conjugated to Alexa Fluor 350  |  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human LARG |  |
|-----------------------|--|--|
| Isotype:              | IgG  |  |
| Cross-Reactivity:     | Mouse  |  |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Pig,Rabbit                             |  |
| Purification:         | Purified by Protein A.                                   |  |

## **Target Details**

| Target:           | ARHGEF12                 |
|-------------------|--------------------------|
| Alternative Name: | LARG (ARHGEF12 Products) |

## Target Details

| Background:         | Synonyms: ARHGEF12, KIAA0382, leukemia-associated rho guanine nucleotide exchange              |  |
|---------------------|--|--|
|                     | factor, Leukemia-associated RhoGEF, PRO2792, Rho guanine exchange factor GEF 12, Rho           |  |
|                     | guanine nucleotide exchange factor GEF 12, Rho guanine nucleotide exchange factor 12,          |  |
|                     | Similar to Mouse Lsc oncogene, ARHGC_HUMAN.  |  |
|                     | Background: Rho GTPases play a fundamental role in numerous cellular processes that are        |  |
|                     | initiated by extracellular stimuli that work through G protein coupled receptors. The encoded  |  |
|                     | protein may form a complex with G proteins and stimulate Rho-dependent signals. This protein   |  |
|                     | is observed to form myeloid/lymphoid fusion partner in acute myeloid leukemia (from            |  |
|                     | EntrezGene). LARG has also been implicated in linking heterotrimeric G proteins to microtubule |  |
|                     | organisation in regulation of cell polarity.   |  |
| Gene ID:            | 23365  |  |
| Pathways:           | Neurotrophin Signaling Pathway, Regulation of G-Protein Coupled Receptor Protein Signaling     |  |
| Application Details |  |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |  |
|                     | IF(IHC-F) 1:50-200   |  |
|                     | IF(ICC) 1:50-200   |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Concentration:      | 1 μg/μL  |  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and    |  |
|                     | 50 % Glycerol.   |  |
| Preservative:       | ProClin  |  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be            |  |
|                     | handled by trained staff only.   |  |
| Storage:            | -20 °C   |  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.              |  |
| Expiry Date:        | 12 months  |  |