

Datasheet for ABIN904920

anti-Lipin 1 antibody (AA 501-600) (AbBy Fluor® 350)



Overview

Overview			
Quantity:	100 μL		
Target:	Lipin 1 (LPIN1)		
Binding Specificity:	AA 501-600		
Reactivity:	Rat		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This Lipin 1 antibody is conjugated to AbBy Fluor® 350		
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))		
Product Details			
Immunogen:	KLH conjugated synthetic peptide derived from human Lipin 1		
Isotype:	IgG		
Cross-Reactivity:	Rat		
Predicted Reactivity:	Human, Mouse, Dog, Cow, Sheep, Pig, Rabbit, Chimpanzee, Goose		
Purification:	Purified by Protein A.		
Target Details			
Target:	Lipin 1 (LPIN1)		
Alternative Name:	Lipin 1 (LPIN1 Products)		

Target Details

Background:	Synonyms: KIAA0188, LPIN1, PAP1, Phosphatidate phosphatase LPIN1, LPIN1_HUMAN.		
	Background: Lipin 1 is a member of the Lipin family of nuclear proteins. This family contains		
	three members: Lipin 1, Lipin 2 and Lipin 3, all of which contain a nuclear signal sequence, a		
	highly conserved amino-terminal (NLIP) domain and a carboxy-terminal (CLIP) domain. LPIN1		
	(Lipin 1) is crucial for normal adipose tissue development and metabolism. LPIN1 selectively		
	activates a subset of PGC1 alpha target pathways, including fatty acid oxidation and		
	mitochondrial oxidative phosphorylation by inducing expression of the nuclear receptor		
	PPARalpha. LPIN1 also inactivates the lipogenic program and suppresses circulating lipid		
	levels. An abundance of LPIN1 promotes fat accumulation and insulin sensitivity, whereas a		
	deficiency in LPIN1 may deter normal adipose tissue development, resulting in insulin		
	resistance and lipodystrophy, a heterogeneous group of disorders characterized by loss of body		
	fat, fatty liver, hypertriglyceridemia and insulin resistance.		
Gene ID:	23175		
Pathways:	Monocarboxylic Acid Catabolic Process		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		

Н	and	lin	C

Expiry Date:

12 months