antibodies .- online.com





anti-Lipin 1 antibody (AA 201-300) (AbBy Fluor® 350)



Go to	D	1	
	Pron	וויאו וו	nane

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	Lipin 1 (LPIN1)
Binding Specificity:	AA 201-300
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Lipin 1 antibody is conjugated to AbBy Fluor® 350
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse Lpin1 protein
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	Lipin 1 (LPIN1)
Alternative Name:	Lpin1 protein (LPIN1 Products)
Background:	Synonyms: fld, Lipin1, Kiaa0188, mKIAA0188, 4631420P06, Phosphatidate phosphatase LPIN1,

Fatty liver	dystronhy	nrotein	I inin-1	I nin1
i accy iivoi	a y o ti o p i i y	protein,	p,	

Background: Plays important roles in controlling the metabolism of fatty acids at differents levels. Acts as a magnesium-dependent phosphatidate phosphatase enzyme which catalyzes the conversion of phosphatidic acid to diacylglycerol during triglyceride, phosphatidylcholine and phosphatidylethanolamine biosynthesis. Acts also as nuclear transcriptional coactivator for PPARGC1A/PPARA regulatory pathway to modulate lipid metabolism gene expression. Is involved in adipocyte differentiation. Isoform 1 is recruited at the mitochondrion outer membrane and is involved in mitochondrial fission by converting phosphatidic acid to diacylglycerol.

Gene ID:	14245
----------	-------

UniProt: Q91ZP3

Pathways: Monocarboxylic Acid Catabolic Process

Application Details

Application Notes: IF(IHC-P) 1:5	50-200
----------------------------------	--------

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months