antibodies -online.com





anti-ZC3H12A antibody (AA 201-300) (Alexa Fluor 350)



_		
()Ver	view	

Quantity:	100 μL
Target:	ZC3H12A
Binding Specificity:	AA 201-300
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZC3H12A antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MCPIP1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ZC3H12A
Alternative Name:	MCPIP1 (ZC3H12A Products)

Target Details

Synonyms: dJ423B22.1, MCP 1 treatment-induced protein, MCP-induced protein 1, MCPIP 1,
MCPIP, Ribonuclease ZC3H12A, RP3-423B22.1, ZC12A_HUMAN, ZC3H 12A, ZC3H12 A,
Zc3h12a, Zinc finger CCCH domain-containing protein 12A, Zinc finger CCCH type containing
12A.
Background: Has RNase activity and selectively degrades specific target mRNA species.
Modulates the immune response and inflammation by regulating the decay of specific mRNA
molecules. Recognizes the 3'-untranslated region (UTR) of the mRNA for IL6, CALCR and IL12B
Required for normal decay of IL6 mRNA (By similarity). Triggers apoptosis and promotes
angiogenesis in response to the binding of CCL2 to CCR2. Regulates expression of CDH12 and CHD19.
80149
Cellular Response to Molecule of Bacterial Origin, Positive Regulation of fat Cell Differentiation
FCM 1:20-100
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.
-20 °C
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
12 months