# antibodies -online.com





## anti-Methionine Adenosyltransferase antibody (AA 31-130) (AbBy Fluor® 488)



Go to Product page

0			
$\cup$	/ 🔾	1 V I	v v

or® 488
munofluorescence

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human Methionine adenosyltransferase	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human, Mouse, Dog, Cow, Pig, Chicken, Rabbit, Zebrafish, Monkey	
Purification:	Purified by Protein A.	

#### **Target Details**

Target:	Methionine Adenosyltransferase (MAT)
Alternative Name:	Methionine adenosyltransferase (MAT Products)

### **Target Details**

9	
Background:	Synonyms: MAT2A AdoMet synthetase 2, AdoMet synthetase, AMS 2, AMS2, MAT 2A, MAT II,
	MATA 2, MATA2, MATII, Methionine adenosyltransferase 2, Methionine adenosyltransferase,
	Methionine adenosyltransferase II alpha, Methionine adenosyltransferase II, S
	adenosyl methion in esynthetasegammaform, Sadenosyl methion in esynthetaseiso formtype
	2, SAMS 2, SAMS2, METK1_HUMAN.
	Background: S adenosylmethionine synthetase catalyzes the formation of S
	adenosylmethionine from methionine and ATP. In mammalian tissues, there are three distinct
	forms of AdoMet synthases designated as alpha, beta, and gamma. Alpha and beta are
	expressed only in adult liver, while gamma is widely distributed in extrahepatic tissues.
Gene ID:	4143
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months