antibodies -online.com





anti-Myosin VIIA antibody (AA 851-950) (Alexa Fluor 555)



Go to Product page

| \sim | | | | | |
|--------|-----|----|-----|-------------|----|
| | 1/6 | ⊃r | \/I | \triangle | ٨/ |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | Myosin VIIA (MYO7A) | |
| Binding Specificity: | AA 851-950 | |
| Reactivity: | Human, Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This Myosin VIIA antibody is conjugated to Alexa Fluor 555 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human Myosin VIIa | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse | |
| Predicted Reactivity: | Rat,Dog,Cow,Pig,Horse,Chicken | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | Myosin VIIA (MYO7A) |
|-------------------|----------------------------|
| Alternative Name: | Myosin 7a (MYO7A Products) |

Target Details

| Background: | Synonyms: DFNB2, MYU7A, NSRD2, USH1B, DFNA11, MYOVIIA, Unconventional myosin-VIIa, |
|---------------------|--|
| | MY07A Real/ground: Museine are actin based mater malegules with ATReas activity. Unconventional |
| | Background: Myosins are actin-based motor molecules with ATPase activity. Unconventional |
| | myosins serve in intracellular movements. Their highly divergent tails bind to membranous |
| | compartments, which are then moved relative to actin filaments. In the retina, plays an |
| | important role in the renewal of the outer photoreceptor disks. Plays an important role in the |
| | distribution and migration of retinal pigment epithelial (RPE) melanosomes and phagosomes, |
| | and in the regulation of opsin transport in retinal photoreceptors. In the inner ear, plays an |
| | important role in differentiation, morphogenesis and organization of cochlear hair cell bundles. |
| | Involved in hair-cell vesicle trafficking of aminoglycosides, which are known to induce |
| | ototoxicity (By similarity). Motor protein that is a part of the functional network formed by |
| | USH1C, USH1G, CDH23 and MY07A that mediates mechanotransduction in cochlear hair cells. |
| | Required for normal hearing. |
| Gene ID: | 4647 |
| UniProt: | Q13402 |
| Pathways: | Sensory Perception of Sound |
| Application Details | |
| Application Notes: | IF(IHC-P) 1:50-200 |
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| | |

Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |