antibodies -online.com





anti-Nerve Growth Factor Receptor (TNFRSF16) Associated Protein 1 (NGFRAP1) (AA 34-111) antibody (Biotin)



Go to Product page

0\	/ (. v .	 νv

Quantity:	100 μL
Target:	Nerve Growth Factor Receptor (TNFRSF16) Associated Protein 1 (NGFRAP1)
Binding Specificity:	AA 34-111
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NADE/NGFRAP1	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Dog,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	Nerve Growth Factor Receptor (TNFRSF16) Associated Protein 1 (NGFRAP1)
Alternative Name:	NGFRAP1 (NGFRAP1 Products)

Target Details

Synonyms: BEX3, BEX3_HUMAN, Brain expressed X linked 3 mouse homolog, Brain-expressed
X-linked protein 3, Nerve growth factor receptor TNFRSF16 associated protein 1, Nerve growth
factor receptor associated protein 1, Nerve growth factor receptor-associated protein 1,
NGFRAP1, Ovarian granulosa cell 13.0 kDa protein HGR74, Ovarian granulosa cell protein,
p75NTR associated cell death executor, p75NTR-associated cell death executor, Protein BEX3.
Background: May be a signaling adapter molecule involved in p75NTR-mediated apoptosis
induced by NGF. Plays a role in zinc-triggered neuronal death (By similarity). May play an
important role in the pathogenesis of neurogenetic diseases. Tissue specificity: Found in
ovarian granulosa cells, testis, prostate and seminal vesicle tissue. High levels also detected in
liver.
27018
Neurotrophin Signaling Pathway, Positive Regulation of Endopeptidase Activity
IHC-P 1:200-400
IHC-F 1:100-500
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.
-20 °C
Store at -20°C for 12 months.