antibodies -online.com







anti-NAT10 antibody (AA 51-150) (Cy5)



\sim				
	11/6	٦r١	/10	۱۸.

Quantity:	100 μL	
Target:	NAT10	
Binding Specificity:	AA 51-150	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NAT10 antibody is conjugated to Cy5	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NAT10
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	NAT10
Alternative Name:	NAT10 (NAT10 Products)
Synonyms: ALP, hALP, KIAA1709, N acetyltransferase 10, N acetyltransferase 10 GCN5 rela	

N acetyltransferase like, N acetyltransferase like protein, N-acetyltransferase 10, NAT10,
NAT10 HUMAN, NET43.

Background: Has protein acetyltransferase activity in vitro. Can acetylate both histones and microtubules. Histone acetylation may regulate transcription and mitotic chromosome decondensation. Activates telomerase activity by stimulating the transcription of TERT, and may also regulate telomerase function by affecting the balance of telomerase subunit assembly, disassembly, and localization. Acetylates alpha-tubulin, which may affect microtubule stability and cell division.

Gene ID:

55226

Application Details

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200

Restrictions:

For Research Use only

IF(ICC) 1:50-200

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months