

Datasheet for ABIN907715 anti-NAT10 antibody (AA 51-150) (Cy7)



Go to Product page

| Overview | |
|-----------------------|---|
| Quantity: | 100 μL |
| Target: | NAT10 |
| Binding Specificity: | AA 51-150 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This NAT10 antibody is conjugated to Cy7 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from human NAT10 |
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | NAT10 |
| Alternative Name: | NAT10 (NAT10 Products) |
| Background: | Synonyms: ALP, hALP, KIAA1709, N acetyltransferase 10, N acetyltransferase 10 GCN5 related, |

| N acetyltransferase like, N acetyltransferase like protein, N-acetyltransferase 10, NAT10, | |
|--|--|
| NAT10 HUMAN, NET43. | |

Background: Has protein acetyltransferase activity in vitro. Can acetylate both histones and microtubules. Histone acetylation may regulate transcription and mitotic chromosome decondensation. Activates telomerase activity by stimulating the transcription of TERT, and may also regulate telomerase function by affecting the balance of telomerase subunit assembly, disassembly, and localization. Acetylates alpha-tubulin, which may affect microtubule stability and cell division.

Gene ID:

55226

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |