

Datasheet for ABIN907862

anti-NDUFV2 antibody (AA 101-200) (AbBy Fluor® 647)



Go to Product page

(۱۱/		٢V	Ĺ		۱٨	١.
	, v	\cup	V	1	$\overline{}$	٧	V

Background:

Quantity: 100 µL Target: NDUFV2 Binding Specificity: AA 101-200 Reactivity: Human Host: Rabbit Clonality: Polyclonal Conjugate: This NDUFV2 antibody is conjugated to AbBy Fluor® 647
Binding Specificity: AA 101-200 Reactivity: Human Host: Rabbit Clonality: Polyclonal
Reactivity: Human Host: Rabbit Clonality: Polyclonal
Host: Rabbit Clonality: Polyclonal
Clonality: Polyclonal
Conjugate: This NDUFV2 antibody is conjugated to AbBy Fluor® 647
Application: Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details
Immunogen: KLH conjugated synthetic peptide derived from human NDUFV2
Isotype: IgG
Predicted Reactivity: Human, Mouse, Rat, Dog, Cow, Pig, Horse, Chicken, Rabbit
Purification: Purified by Protein A.
Target Details
Target: NDUFV2
Alternative Name: NDUFV2 (NDUFV2 Products)

Synonyms: 24 kDa subunit of Complex I, CI-24k, NADH dehydrogenase ubiquinone flavoprotein

2, 24 kDa, NADH ubiquinone oxidoreductase 24 kDa subunit, Ubiquinoneflavoprotein 2, mitochondrial precursor.

Background: The NADH-ubiquinone oxidoreductase complex (complex I) of the mitochondrial respiratory chain catalyzes the transfer of electrons from NADH to ubiquinone, and consists of at least 43 subunits. The complex is located in the inner mitochondrial membrane. This gene encodes the 24 kDa subunit of complex I, and is involved in electron transfer. Mutations in this gene are implicated in Parkinson's disease, bipolar disorder, schizophrenia, and have been found in one case of early onset hypertrophic cardiomyopathy and encephalopathy. A non-transcribed pseudogene of this locus is found on chromosome 19. [provided by RefSeq, Oct 2009].

Gene ID:

4729

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months