## antibodies .- online.com







## anti-RTN4R antibody (AA 151-350) (Alexa Fluor 555)



| ( ) | 1/0 | r\ /1 | 014 |   |
|-----|-----|-------|-----|---|
| ( ) | ve  | I V I | -v  | V |
|     |     |       |     |   |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | RTN4R  |
| Binding Specificity: | AA 151-350   |
| Reactivity:          | Mouse, Rat   |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This RTN4R antibody is conjugated to Alexa Fluor 555   |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## **Product Details**

| Immunogen:        | KLH conjugated synthetic peptide derived from mouse Nogo Receptor |  |
|-------------------|---|--|
| Isotype:          | IgG   |  |
| Cross-Reactivity: | Mouse, Rat  |  |
| Purification:     | Purified by Protein A.  |  |

## **Target Details**

| Target:   | RTN4R                          |
|---|--------------------------------|
| Alternative Name:   | Nogo Receptor (RTN4R Products) |
| Background: Synonyms: NgR, NgR1, NOGOR, Reticulon-4 receptor, Nogo receptor, Nogo-66 receptor, Rt |                                |

| Target Details      |   |
|---------------------|---|
|                     | Background: Receptor for RTN4, OMG and MAG. Mediates axonal growth inhibition and may play a role in regulating axonal regeneration and plasticity in the adult central nervous system (By similarity). Acts in conjunction with RTN4 and LIGO1 in regulating neuronal precursor cell motility during cortical development. |
| Gene ID:            | 65079   |
| UniProt:            | Q99PI8  |
| Pathways:           | Neurotrophin Signaling Pathway  |
| Application Details |   |
| Application Notes:  | IF(IHC-P) 1:50-200<br>IF(IHC-F) 1:50-200<br>IF(ICC) 1:50-200  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 μg/μL   |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be   |

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

handled by trained staff only.

-20 °C

12 months

Storage:

Expiry Date:

Storage Comment: