# antibodies -online.com





# anti-NSP5 antibody (AA 601-700) (Cy3)



Go to Product page

| ( ) | 11/0   | K\ / | iew                 | 1 |
|-----|--------|------|---------------------|---|
|     | $\cup$ | ועוי | $\square \vee \vee$ | ı |
|     |        |      |                     |   |

| Quantity:            | 100 μL   |
|----------------------|--|
| Target:              | NSP5 (SPECC1)  |
| Binding Specificity: | AA 601-700   |
| Reactivity:          | Rat  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This NSP5 antibody is conjugated to Cy3  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human NSP5 |
|-----------------------|--|
| Isotype:              | IgG  |
| Cross-Reactivity:     | Rat  |
| Predicted Reactivity: | Human,Dog,Sheep,Rabbit                                   |
| Purification:         | Purified by Protein A.                                   |

## **Target Details**

| Target:           | NSP5 (SPECC1)          |
|-------------------|------------------------|
| Alternative Name: | NSP5 (SPECC1 Products) |

## **Target Details**

| Synonyms: Cytokinesis and spindle organization B, Cytospin B, FLJ36955, HCMOGT 1, NSP,                |  |
|---|--|
| NSP5, Nuclear structure protein 5, SPECC1, Spectrin domain with coiled coils 1, Sperm antigen         |  |
| HCMOGT 1, Sperm antigen with calponin homology and coiled coil domains 1, Structure protein           |  |
| NSP5a3a, Structure protein NSP5a3b, Structure protein NSP5b3a, Structure protein NSP5b3b,             |  |
| CYTSB_HUMAN.  |  |
| Background: NSP type 5 alpha 3 alpha is a novel structure protein belonging to the cytospin-A         |  |
| family, that may play a role in nuclear function during cell division. It is also highly expressed in |  |
| certain tumor cell lines, suggesting that it may serve as a good tumor marker. It was was             |  |
| cloned as part of a chromosomal translocation involving PDGFRB in a case of juvenile                  |  |
| myelomonocytic leukemia.  |  |
| 92521   |  |
|   |  |
| IF(IHC-P) 1:50-200  |  |
| IF(IHC-F) 1:50-200  |  |
| IF(ICC) 1:50-200  |  |
| For Research Use only   |  |
|   |  |
| Liquid  |  |
| 1 μg/μL   |  |
| Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and           |  |
| 50 % Glycerol.  |  |
| ProClin   |  |
| This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be                   |  |
| handled by trained staff only.  |  |
| -20 °C  |  |
| Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                     |  |
| 12 months   |  |
|   |  |