# antibodies -online.com





# anti-PPAP2B antibody (AA 241-311) (Alexa Fluor 350)



Go to Product page

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL	
Target:	PPAP2B	
Binding Specificity:	AA 241-311	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PPAP2B antibody is conjugated to Alexa Fluor 350	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human PAP2B
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

## **Target Details**

Target:	PPAP2B	
Alternative Name:	PAP2B (PPAP2B Products)	

### **Target Details**

Synonyms: LPP3, VCIP, Dri42, PAP2B, Lipid phosphate phosphohydrolase 3, PAP2-beta,	
Phosphatidate phosphohydrolase type 2b, Phosphatidic acid phosphatase 2b, PAP-2b, Vascular	
endothelial growth factor and type I collagen-inducible protein, PPAP2B	
Background: Catalyzes the conversion of phosphatidic acid (PA) to diacylglycerol (DG). In	
addition it hydrolyzes lysophosphatidic acid (LPA), ceramide-1-phosphate (C-1-P) and	
sphingosine-1-phosphate (S-1-P). The relative catalytic efficiency is LPA = PA > C-1-P > S-1-P.	
May be involved in cell adhesion and in cell-cell interactions.	
8613	
014495	
IF(IHC-P) 1:50-200	
IF(IHC-F) 1:50-200	
IF(ICC) 1:50-200	
For Research Use only	
Liquid	
1 μg/μL	
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
ProClin	
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
handled by trained staff only.	
-20 °C	
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
12 months	