antibodies .- online.com





anti-PDGFA antibody (AA 125-170) (Alexa Fluor 350)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	ועוי	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	PDGFA	
Binding Specificity:	AA 125-170	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PDGFA antibody is conjugated to Alexa Fluor 350	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PDGF-A	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Predicted Reactivity:	Rat,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	PDGFA
Alternative Name:	PDGF A (PDGFA Products)

Target Details

rarget Details		
Background:	Synonyms: PDGF1, PDGF-A, Platelet-derived growth factor subunit A, PDGF subunit A, PDGF-1,	
	Platelet-derived growth factor A chain, Platelet-derived growth factor alpha polypeptide, PDGFA	
	Background: Growth factor that plays an essential role in the regulation of embryonic	
	development, cell proliferation, cell migration, survival and chemotaxis. Potent mitogen for cells	
	of mesenchymal origin. Required for normal lung alveolar septum formation during	
	embryogenesis, normal development of the gastrointestinal tract, normal development of	
	Leydig cells and spermatogenesis. Required for normal oligodendrocyte development and	
	normal myelination in the spinal cord and cerebellum. Plays an important role in wound healing.	
	Signaling is modulated by the formation of heterodimers with PDGFB (By similarity).	
Gene ID:	5154	
UniProt:	P04085	
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin	
	Signaling Pathway, Smooth Muscle Cell Migration, Platelet-derived growth Factor Receptor	
	Signaling	
A 1: 1: D 1:1		
Application Details		
Application Notes:	FCM 1:20-100	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months