antibodies .- online.com







anti-PPARGC1A antibody (AA 151-250) (Cy7)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	PPARGC1A
Binding Specificity:	AA 151-250
Reactivity:	Human, Rat, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PPARGC1A antibody is conjugated to Cy7
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PGC1 alpha
Isotype:	IgG
Cross-Reactivity:	Human, Pig, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	PPARGC1A
Alternative Name:	PGC1 alpha (PPARGC1A Products)

Background:

Synonyms: LEM6, Ligand effect modulator 6, PERC, Peroxisome prolerative activated receptor, gamma, coactivator 1 alpha, Peroxisome prolerative activated receptor, gamma, coactivator 1, Peroxisome prolerator activated receptor gamma coactivator 1 alpha, peroxisome prolerator-activated receptor gamma coactivator 1 beta, Peroxisome prolerator-activated receptor gamma coactivator 1-alpha, Peroxisome prolerator-activated receptor gamma coactivator 1-beta, peroxisome prolerator-activated receptor gamma, coactivator 1 beta, PGC 1 alpha, PGC 1 alpha, PGC 1v, PGC-1beta, PGC-1-alpha, PGC-1-beta, PGC-1-related estrogen receptor alpha coactivator, PGC1, PGC1alpha, PGC1A, PGC1v, PPAR gamma coactivator 1 alpha, PPAR gamma coactivator 1 alpha 3 ligand effect modulator 6, PPAR gamma coactivator 1, PPAR gamma coactivator variant form, PPAR gamma coactivator-1beta, PPAR-gamma coactivator 1-alpha, PPAR-gamma coactivator 1-beta, PPARGC-1-alpha, PPARGC-1-beta, PPARGC-1-beta, PPARGC1, PPARGC1A, Ppargc1b, PRGC1_HUMAN, PRGC2_HUMAN.

Background: The protein encoded by this gene is a transcriptional coactivator that regulates the genes involved in energy metabolism. This protein interacts with PPARgamma, which permits the interaction of this protein with multiple transcription factors. This protein can interact with, and regulate the activities of, cAMP response element binding protein (CREB) and nuclear respiratory factors (NRFs). It provides a direct link between external physiological stimuli and the regulation of mitochondrial biogenesis, and is a major factor that regulates muscle fiber type determination. This protein may be also involved in controlling blood pressure, regulating cellular cholesterol homoeostasis, and the development of obesity. [provided by RefSeq].

Molecular Weight: 91-110kDa

Gene ID: 10891

UniProt: Q9UBK2

Pathways: Intracellular Steroid Hormone Receptor Signaling Pathway, Carbohydrate Homeostasis,

Intracellular Steroid Hormone Receptor Signaling Pathway, Carbohydrate Homeostasis,
Regulation of Lipid Metabolism by PPARalpha, Regulation of Carbohydrate Metabolic Process,
Smooth Muscle Cell Migration, Brown Fat Cell Differentiation

Application Details

Application Notes: FCM 1:20-100

IF(IHC-P) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months