

Datasheet for ABIN911300 anti-PKMYT1 antibody (AA 121-220) (AbBy Fluor® 350)



Overview	
Quantity:	100 µL
Target:	PKMYT1
Binding Specificity:	AA 121-220
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PKMYT1 antibody is conjugated to AbBy Fluor® 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human PKMYT1
Isotype:	lgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Horse
Purification:	Purified by Protein A.
Target Details	
Target:	PKMYT1
Alternative Name:	PKMYT1 (PKMYT1 Products)
Background:	Synonyms: cdc2 inhibitory kinase, Membrane associated tyrosine and threonine specic cdc2

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN911300 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

	inhibitory kinase, Membrane-associated tyrosine- and threonine-specic cdc2-inhibitory kinase,
	MYT1, Myt1 kinase, PKMYT 1, Pkmyt1, PMYT1_HUMAN, Protein kinase membrane associated
	tyrosine/threonine 1, Protein kinase Myt1, DKFZp547K1610, FLJ20093.
	Background: PKMYT1 is a member of the serine/threonine protein kinase family. It
	preferentially phosphorylates and inactivates cell division cycle 2 protein (cdc2), and thus acts
	as a negative regulator of entry into mitosis (G2 to M transition). It mediates phosphorylation of
	cdc2 predominantly on 'Thr-14' and is also involved in Golgi fragmentation. It may be involved in
	phosphorylation of cdc2 on 'Tyr-15' to a lesser degree, however tyrosine kinase activity is
	unclear and may be indirect. It may be a downstream target of Notch signaling pathway during
	eye development. PKMYT1 is negatively regulated by hyperphosphorylation during mitosis.
Gene ID:	9088
Gene ID: Pathways:	9088 Mitotic G1-G1/S Phases, M Phase
Pathways:	
Pathways: Application Details	Mitotic G1-G1/S Phases, M Phase
Pathways: Application Details	Mitotic G1-G1/S Phases, M Phase IF(IHC-P) 1:50-200

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months