antibodies .- online.com





anti-PLCL1 antibody (AA 1001-1096) (Alexa Fluor 555)



and the following the first term of the first te

()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	ı١	νı	ㄷ	٧	۷

Quantity:	100 μL
Target:	PLCL1
Binding Specificity:	AA 1001-1096
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PLCL1 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PLCL1	
Isotype:	IgG	
Specificity:	This antibody may exhibit a slight secondary cross-reactivity with PLCL2 due to a 69 % non-sequential sequence similarity over the immunogen range.	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse	
Purification:	Purified by Protein A.	

Target Details

Target:	PLCL1
Alternative Name:	PLCL1 (PLCL1 Products)
Background:	Synonyms: PLCL1, PLCE, Phospholipase C deleted in lung carcinoma, Phospholipase C epsilon,
	Phospholipase C like 1, Phospholipase C related but catalytically inactive protein,
	Phospholipase C related, but catalytically inactive protein, PLCE, PLCL, PRIP, Plcl1,
	C230017K02Rik, KIAA1092, PLC-L, PLC-L1, PLCE, PLCL, PRIP, PRIP-1, PLCL1_HUMAN.
	Background: PLCL1 is involved an inositol phospholipid based intracellular signaling cascade.
	Shows no PLC activity to phosphatidylinositol 4,5-bisphosphate and sphatidylinositol.
	Component in the dependent endocytosis process of GABA A receptor (By similarity).
	Regulates the turnover of receptors and thus contributes to the maintenance of GABA
	mediated synaptic inhibition. Its aberrant expression could contribute to the genesis and
	progression of lung carcinoma. Acts as an inhibitor of PPP1C.
Gene ID:	5334
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months