antibodies -online.com







anti-Relaxin 1 antibody (AA 51-150) (Alexa Fluor 555)



Go to Product page

| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | Relaxin 1 (RLN1) | |
| Binding Specificity: | AA 51-150 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This Relaxin 1 antibody is conjugated to Alexa Fluor 555 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human Relaxin 1 |
|-----------------------|---|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat |
| Purification: | Purified by Protein A. |

Target Details

| Target: | Relaxin 1 (RLN1) |
|---|---------------------------|
| Alternative Name: | Relaxin 1 (RLN1 Products) |
| Background: Synonyms: H1 antibody Preprorelaxin H1, PreprorelaxinH1, Prorelaxin, Prorelaxin H1, | |

ProrelaxinH1, Relaxin 1 H1, Relaxin H1, Relaxin1, Relaxin-1, RelaxinH1, RLN 1, RLN1, RLX H1, RLXH1, RLXH1, REL1_HUMAN.

Background: Relaxin is a peptide hormone produced by the corpora lutea of ovaries during pregnancy in many mammalian species, including man. The secretion of the hormone into the blood stream just before parturition results in a marked softening and lengthening of the pubic symphysis and a softening of the cervix, which facilitates the birth process. By inhibiting uterine contractions, relaxin may influence the timing of parturition. Like insulin, relaxin consists of two peptide chains, A and B, covalently linked by disulfide bonds. By further analogy to insulin, the two peptides are synthesized as a single-chain precursor polypeptide with the B chain at the NH2-terminus.

Gene ID:

6013

Application Details

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200 IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

| Format: | Liquid | |
|--------------------|--|--|
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |