antibodies -online.com





anti-RELM alpha antibody (AA 51-111) (Alexa Fluor 350)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Background:

Overview	
Quantity:	100 μL
Target:	RELM alpha
Binding Specificity:	AA 51-111
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RELM alpha antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc))
Product Details	
Product Details Immunogen:	KLH conjugated synthetic peptide derived from mouse RELMa
	KLH conjugated synthetic peptide derived from mouse RELMa
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	IgG Mouse, Rat
Immunogen: Isotype: Cross-Reactivity: Purification:	IgG Mouse, Rat

Synonyms: HIMF, Xcp2, Fizz1, RELMa, Fizz-1, RELMalpha, RELM-alpha, 18119L16Rik, Resistin-

like alpha, Cysteine-rich secreted protein A12-gamma, Cysteine-rich secreted protein FIZZ1,

Hypoxia-induced mitogenic factor, Parasite-induced macrophage novel gene 1 protein, Retnla, Pmng1

Background: A family of resistin-like molecules (RELMs) has been identified in rodents and humans. RELM alpha belongs to a unique family of tissue-specific cytokines termed FIZZ (found in inflammatory zone) and RELM. The three known members of this family, Resistin, RELM alpha and RELM beta are 85-94 amino acid secreted proteins sharing a conserved C-terminal domain. RELM alpha and Resistin are secreted exclusively by adipocytes while RELM beta is expressed in the epithelium of the colon and small bowel. The RELMs together with resistin comprise a class of tissue-specific signaling molecules. The physiological role and molecular targets of RELM alpha are still unknown.

Gene ID:

57262

UniProt:

Q9EP95

Application Details

Application Notes: IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	