.-online.com antibodies

Datasheet for ABIN913845 anti-Cbl Proto-Oncogene B, E3 Ubiquitin Protein Ligase (CBLB) (AA 101-200) antibody (Alexa Fluor 350)



Overview

Quantity:	100 µL	
Target:	Cbl Proto-Oncogene B, E3 Ubiquitin Protein Ligase (CBLB)	
Binding Specificity:	AA 101-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	Alexa Fluor 350	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RNF56/CBL-B
lsotype:	lgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Pig,Horse,Chicken
Purification:	Purified by Protein A.
Target Details	
Target:	Cbl Proto-Oncogene B, E3 Ubiquitin Protein Ligase (CBLB)
Alternative Name:	RNF56 (CBLB Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN913845 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Background:	Synonyms: Cbl-b, RNF56, Nbla00127, E3 ubiquitin-protein ligase CBL-B, Casitas B-lineage	
Backyrounu.	lymphoma proto-oncogene b, RING finger protein 56, SH3-binding protein CBL-B, Signal	
	transduction protein CBL-B, CBLB	
	Background: E3 ubiquitin-protein ligase which accepts ubiquitin from specific E2 ubiquitin-	
	conjugating enzymes, and transfers it to substrates, generally promoting their degradation by	
	the proteasome. Negatively regulates TCR (T-cell receptor), BCR (B-cell receptor) and FCER1	
	(high affinity immunoglobulin epsilon receptor) signal transduction pathways. In naive T-cells,	
	inhibits VAV1 activation upon TCR engagement and imposes a requirement for CD28	
	costimulation for proliferation and IL-2 production. Also acts by promoting PIK3R1/p85	
	ubiquitination, which impairs its recruitment to the TCR and subsequent activation. In activated	
	T-cells, inhibits PLCG1 activation and calcium mobilization upon restimulation and promotes	
	anergy. In B-cells, acts by ubiquitinating SYK and promoting its proteasomal degradation.	
	Slightly promotes SRC ubiquitination. May be involved in EGFR ubiquitination and	
	internalization. May be functionally coupled with the E2 ubiquitin-protein ligase UB2D3.	
Gene ID:	868	
UniProt:	Q13191	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 µg/µL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN913845 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

lond	lina
land	
10110	

Storage Comment:

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN913845 | 03/07/2024 | Copyright antibodies-online. All rights reserved.